Sickness Policy

If a child is considered to be unwell while attending the setting, a senior member of staff will assess the child. If a child is unwell, we will contact the parent/carer, or one of the emergency contacts, for the child to be collected.

In emergencies we will follow our first aid procedures and phone for paramedics. Where appropriate we will administer medicine or medical care in accordance with our Medication policy.

If a child is unwell, or has an infectious illness we require them to be kept at home until the infectious period is over. The Department of Health recommend the following periods for children to be kept away from nursery:

Illness		-
Chicken pox At least 5 days from onset of rash and until all blisters have crusted over Conjunctivitis * Following Doctor's appointment: Bacterial - 24 hours from onset, with medication. Viral - no exclusion with letter from doctor Respiratory infections including coronavirus (COVID-19) If they have a high temperature and are unwell children should not attend until better If positive test result for COVID-19 they should not attend 3 days after test Diarrhoea/ Vomiting Flu (influenza) Until recovered Hepititis A T days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough At least 5 days from commencing antibiotics or 21 days without antibiotic	Illness	Exclusion Period
Conjunctivitis * Following Doctor's appointment: Bacterial - 24 hours from onset, with medication. Viral - no exclusion with letter from doctor If they have a high temperature and are unwell children should not attend until better If positive test result for COVID-19 they should not attend 3 days after test Diarrhoea/ Vomiting Flu (influenza) Until recovered Hepititis A 7 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after on exclusion with letter from onset, with medication. Viral - no exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	Antibiotics prescribed	
Conjunctivitis * Following Doctor's appointment: Bacterial - 24 hours from onset, with medication. Viral - no exclusion with letter from doctor Respiratory infections including coronavirus (COVID-19) If they have a high temperature and are unwell children should not attend until better If positive test result for COVID-19 they should not attend 3 days after test Diarrhoea/ Vomiting	Chicken pox	
with medication. Viral - no exclusion with letter from doctor Respiratory infections including coronavirus (COVID-19) If they have a high temperature and are unwell children should not attend until better If positive test result for COVID-19 they should not attend 3 days after test Diarrhoea/ Vomiting 48 hours diarrhoea &/or vomiting has stopped Flu (influenza) Until recovered Hepititis A 7 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic		
Respiratory infections including coronavirus (COVID-19) (COVID-19) Diarrhoea/ Vomiting Flu (influenza) Hepititis A To days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Impetigo Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps Ringworm Rubella (German measles) Scabies Scarlet Fever Scarlet Fever Schingles Only exclude if rash is weeping and cannot be covered. At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection If they have a high temperature and are unwell children should not attend are unwell children should not attend antibletter If positive test result for COVID-19 they should not attend 3 days after test If positive test result for COVID-19 they should not attend 3 days after test 4 days after test 4 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Until recovered 4 days from onset of rash if well enough Until recovered 5 days from onset of swelling Ringworm Until Treated 5 days from onset of rash 5 days from onset of rash 5 days from onset of rash 6 days from onset of rash 7 days after symptom onset if no jaundice (or 7 days after symptom onset if no jaundice) Until recovered 1 days from onset of rash if well enough Until recovered 5 days from onset of swelling At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough	Conjunctivitis *	
including coronavirus (COVID-19) If positive test result for COVID-19 they should not attend 3 days after test Diarrhoea/ Vomiting		
COVID-19 If positive test result for COVID-19 they should not attend 3 days after test		
after test Diarrhoea/ Vomiting		
Diarrhoea/ Vomiting Flu (influenza) Until recovered Hepititis A 7 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	(COVID-19)	·
Flu (influenza) Hepititis A 7 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Rubella (German measles) Scabies Until Treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough		
Hepititis A 7 days after onset of jaundice (or 7 days after symptom onset if no jaundice) Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic		
Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Until recovered Septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) Whooping cough Total Infection Nealed or 48 hours after starting antibiotic reatment Ferrom onset of rash is well enough Until recovered Stays from onset of swelling Until Treated Stays from onset of rash scapital recommencing antibiotics At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough		
Impetigo Until lesions are crusted or healed or 48 hours after starting antibiotic treatment Measles 4 days from onset of rash if well enough Until recovered Septicaemia* Mumps 5 days from onset of swelling Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) Consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough	Hepititis A	· · · · · · · · · · · · · · · · · · ·
Adays from onset of rash if well enough Meningitis or septicaemia* Mumps		
Measles4 days from onset of rash if well enoughMeningitis or septicaemia*Until recoveredMumps5 days from onset of swellingRingwormUntil TreatedRubella (German measles)5 days from onset of rashScabiesUntil treatedScarlet Fever24 hours after commencing antibioticsShinglesOnly exclude if rash is weeping and cannot be covered.Tuberculosis (pulmonary)At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infectionWhooping cough5 days from commencing antibiotics or 21 days without antibiotic	Impetigo	
Meningitis or septicaemia* Mumps 5 days from onset of swelling Ringworm Until Treated Rubella (German measles) Scabies Until treated Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic		
septicaemia*5 days from onset of swellingRingwormUntil TreatedRubella (German measles)5 days from onset of rashScabiesUntil treatedScarlet Fever24 hours after commencing antibioticsShinglesOnly exclude if rash is weeping and cannot be covered.TuberculosisAt least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infectionWhooping cough5 days from commencing antibiotics or 21 days without antibiotic		
Mumps5 days from onset of swellingRingwormUntil TreatedRubella (German measles)5 days from onset of rashScabiesUntil treatedScarlet Fever24 hours after commencing antibioticsShinglesOnly exclude if rash is weeping and cannot be covered.TuberculosisAt least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infectionWhooping cough5 days from commencing antibiotics or 21 days without antibiotic		Until recovered
Ringworm Rubella (German 5 days from onset of rash 5 days from onset of rash 5 days from onset of rash 6 days from onset of rash 7 days from onset of rash 7 days from onset of rash 8 days from commencing antibiotics 6 days without antibiotic 8 days from commencing antibiotics or 21 days without antibiotic 8 days from commencing antibiotics or 21 days without antibiotic		
Rubella (German measles) Scabies Until treated Scarlet Fever Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) Whooping cough 5 days from onset of rash Onset of rash Stables Until treated 24 hours after commencing antibiotics Only exclude if rash is weeping and cannot be covered. At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough	•	
measles)Until treatedScarlet Fever24 hours after commencing antibioticsShinglesOnly exclude if rash is weeping and cannot be covered.TuberculosisAt least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infectionWhooping cough5 days from commencing antibiotics or 21 days without antibiotic	Ringworm	
Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	`	5 days from onset of rash
Scarlet Fever 24 hours after commencing antibiotics Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	,	
Shingles Only exclude if rash is weeping and cannot be covered. Tuberculosis (pulmonary) Consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough Only exclude if rash is weeping and cannot be covered. At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection 5 days from commencing antibiotics or 21 days without antibiotic		
Tuberculosis (pulmonary) At least 2 weeks after start of effective antibiotic treatment & consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	Scarlet Fever	
 (pulmonary) Consult with local HPA. No exclusion for non-pulmonary or latent TB infection Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic 		
Whooping coughIatent TB infection5 days from commencing antibiotics or 21 days without antibiotic	Tuberculosis	
Whooping cough 5 days from commencing antibiotics or 21 days without antibiotic	(pulmonary)	
Antibiotics 24 hours after commencing antibiotics		
	Antibiotics	24 hours after commencing antibiotics

Where more than 4 children have conjunctivitis then the cases are not isolated and the setting will request that children with viral and bacterial conjunctivitis remain at home in order to prevent an epidemic.

Confidentiality will be preserved, but in cases of infectious illness, other parent/carer's will be notified. Information will include details of the illness, possible symptoms and guidance. In the case of a serious illness, sickness outbreak or notifiable disease (www.hpa.org.uk)

OfSTED and Public Health England (0844 225 3560) will be notified as appropriate. Health Protection Agency will be contacted in any case where further advice is required.

Reviewed: May 2022